





- Bastien P. Comparison of six PCR methods using peripheral blood for detection of canine visceral leishmaniasis J. Clin. Microbiol. 2002;40:210–215.
- [5] Marfurt J, Nasereddin A, Niederwieser I, Jaffe C.L, Beck H.P, Felger I. Identification and differentiation of *Leishmania* species in clinical samples by PCR amplification of the miniexon sequence and subsequent restriction fragment length polymorphism analysis. J. Clin. Microbiol. 2003; 41: 3147–3153.
- [6] Wilson ME, Young BM, Andersen KP, Weinstock JV, Metwali A, Ali KM, *et al.* A recombinant *Leishmania chagasi* antigen that stimulates cellular immune responses in infected mice. Infect. Immun. 1995; 63(5): 2062–2069.
- [7] Kumar J, Bumb R, Ansari N, Mehta R, Salotra P. Cutaneous Leishmaniasis caused by leishmania tropica in BIKANER, INDIA: Parasite identification and characterization using molecular and immunologic tools. Raam J. Trop. Med. Hyg. 2007; 76(5): 896–901.
- [8] Mahdy M.A.K, Al-Mekhlafi H.M, Al-Mekhlafi A.M, Lim Y.A.L, Bin Shuaib N.O.M, Azazy A.A, *et al.* Molecular Characterization of Leishmania Species Isolated from Cutaneous Leishmaniasis in Yemen. PLoS ONE. 2010; 9(5): 12879.
- [9] Farahmand M, Nahrevanian H, Atashi Shirazi H, Naeimi S, Farzanehnejad Z. An overview of a diagnostic and epidemiologic reappraisal of cutaneous leishmaniasis in Iran. Braz J Infect Dis. 2011; 15(1):17-21.
- [10] Singh B, Sundar S. Leishmaniasis: Vaccine candidates and perspectives. Vaccine. 2012; 30: 3834-3842.
- [11] Tabbara K.S. Progress toward leishmania vaccine. Saudi med j. 2006; 27(7): 942-950. [12]. Khamesipour A, Rafati S, Davoudi N, Maboudi F, Modabber F. Leishmaniasis vaccine candidates for development: A global Overview. Indian J Med Res. 2006; 123: 423-438.