





















- [29] E. Portis, A. Acquadro, C. Comino, and S. Lanteri, "Effect of farmer's seed selection on genetic variation of landrace population of pepper (*Capsicum annuum* L.), grown in North-west Italy," *Genet. Res. Crop Evol.*, vol. 51, pp. 581-590, 2004.
- [30] T. J. Kang, and M. S. Yang, "Rapid and reliable extraction of genomic DNA from various wild-type and transgenic," *BMC Biotechnol.*, vol. 4, no. 20, pp. 1-12, 2004.
- [31] S. Sensoy, H. Aktas, and K. Abak, "Genetic diversity in some Turkish pepper (*Capsicum annuum* L.) genotypes revealed by AFLP analyses," *African J. Biotechnol.*, vol. 8, no. 18, pp. 4378-4386, 2009.
- [32] Ø. Hammer, D. A.T. Harper, and P. D. Ryan, "PAST: Palaeontological statistics software package for education and data analysis," *Paleontologia Eletronica*, vol. 4, no. 1, pp. 1-9, 2001.
- [33] E. Hiroaki, I. Hiroyuki, T. Tadashi, and I. Shigera, "Genetic diversity of the peruvianum complex (*Lycopersicon peruvianum* L. Mill and *Lycopersicon chilense* Dun.) revealed by RAPD analysis," *Euphytica*, vol. 116, pp. 23-31, 2000.
- [34] H. P. Young, A. L. Marilyn, M. A. West, and D. A. Stclair, "Evaluation of AFLPs for germplasm finger printing and assessment of genetic diversity in cultivars of tomato (*Lycopersicon esculentum*)," *Genome*, vol. 47, pp. 510-518, 2004.
- [35] S. Garcia-Martinez, L. Andreani, M. Garcia- Gusano, F. Geuna, and J. J. Ruiz, "Evaluation of Amplified fragment length polymorphism and simple sequence repeats for tomato germplasm finger printing: utility for grouping together closely related traditional cultivars," *Genome*, vol. 49, pp. 648-656, 2006.
- [36] S. Lanteri, A. Acquadro, L. Quagliotti, and E. Portis, "RAPD and AFLP assessment of genetic variation in a landrace of pepper (*Capsicum annuum* L.), grown in North-west Italy," *Genet. Res. Crop Evol.*, vol. 50, pp. 723-735, 2003.
- [37] M., Vuylsteke, J. D. Pelema, and M. J. T. VanEijk, "Technology for DNA finger printing," *Nature protocols*, vol. 2, pp. 1387-1398, 2007.
- [38] M. K. Bozokalfa, D. Esiyok, and K. Turhan, "Patterns of phenotypic variation in a germplasm collection of pepper (*Capsicum annuum* L.) from Turkey," *Spanish J. Agri. Res.*, vol. 7, no. 1, pp. 83-95, 2009.
- [39] E. A. Snigir, O. N. Pyshnaya, E. Z. Kochieva, and N. N. Ryzhova, "AFLP-analysis of varietal polymorphism in *Capsicum annuum* L.," *Selskokhozyaistvennaya Biologiya (Agricultural Biology)*, vol. 1, pp. 53-60, 2013.
- [40] S. L. Krishnamurthy, Y. Prashanth, A. M. Rao, K. M. Reddy, and R. Ramachandra, "Assessment of AFLP marker based genetic diversity in chilli (*Capsicum annuum* L. & *C. baccatum* L.)," *Indian J. Biotech.*, Vol. 14, pp. 49-54, 2015.
- [41] J. Baral, and P. W. Bosland, "Genetic diversity of *capsicum* germ plasm collection from Nepal as determined by randomly amplified polymorphic DNAmarkers," *J. Am. Soc. Hortic. Sci.*, vol. 127, pp. 316-324, 2002.
- [42] S. T. Thul, R. K. Lal, A. K. Shasany, M. P. Darokar, A. K. Gupta, M. M. Gupta, R. K. Verma, and S. P. S Khanuja, "Estimation of phenotypic divergence in a collection of *Capsicum* specie for yield related traits," *Euphytica*, vol. 168, pp. 189-196, 2009.
- [43] S. M. Tam, C. Mhiri, A. Vogelaar, M. Kerkveld, S. R. Pearce, and M. A. Grand Bastien, "Comparative analyses of genetic diversities within tomato and pepper collections detected by ret transposon- based SSAP, AFLP and SSR," *Theor. Appl. Genet.*, vol. 110, pp. 819-831, 2005.
- [44] Y. Wahyuni, A. R. Ballester, Y. Tikunov, R. C. H. Devos, K. T. B. Pelgrom, A. Maharijaya, E. Sudarmonowati, R. J. Bino, and A. G. Bovy, "Metabolomics and molecular marker analysis to explore pepper (*Capsicum* sp.) biodiversity," *Metabolomics*, vol. 9, no. 1, pp. 130-144, 2013.
- [45] R. Vellve, "The decline of diversity in European agriculture," *Ecologist*, vol. 23, pp. 64-69, 1993.
- [46] V., Lefebvre, A. Palloix, and M. Rives, "Nuclear RFLP between pepper cultivars (*Capsicum annuum* L.)," *Euphytica*, vol. 71, pp. 189-199, 1993.
- [47] V. Lefebvre, B. Goffinet, J. C. Chauvet, B. Caromel, P. Signoret, R. Brand, and A. Palloix, "Evaluation of genetic distances between pepper inbred lines for cultivar protection purposes: comparison of AFLP, RAPD and phenotypic data," *Theor. Appl. Genet.*, vol. 102, pp. 741-750, 2001.